

ALTAMONTE PEDIATRIC ASSOCIATES

Well Child Care at 6 Months

Feeding - If you haven't started your baby on baby foods (other than cereal), you can start now. Begin with fruits and vegetables. Start one new food at a time for a few days to make sure your baby digests it well. Do not start meats until your baby is 7 to 8 months old. Do not give foods that require chewing. At meals give the baby formula, or breast-feed your baby before giving baby food.

Your baby should continue having breast milk or infant formula until he is 1 year old. Your baby may soon be ready for a cup although it will be messy at first. Try giving a cup occasionally to see if your baby likes it. Don't let him lie down with a bottle. This can lead to tooth decay or ear infections.

Make cereal with formula or breast milk only. Use a spoon to feed your baby cereal, not a bottle or an infant feeder. Sitting up while eating helps your baby learn good eating habits.

Development - At this age babies are usually rolling over and beginning to sit by themselves. Babies squeal, babble, laugh, and often cry very loudly. They may be afraid of people they do not know. Meet your baby's needs quickly and be patient with your baby.

Sleep - 6-month-olds may not want to be put in bed. A favorite blanket or stuffed animal may make bedtime easier. Do not put a bottle in the bed with your baby. Develop a bedtime routine like playing a game, singing a lullaby, turning the lights out, and giving a goodnight kiss. Make the routine the same every night. Be calm and consistent with your baby at bedtime. If your baby is not sleeping through the night, ask your doctor for further information about preventing sleep problems.

Safety Tips

- Avoid Choking and Suffocation**
- Cords, ropes, or strings around the baby's neck can choke him. Keep cords away from the crib.
 - Keep all small, hard objects out of reach.
 - Use only unbreakable toys without sharp edges or small parts that can come loose.
 - Avoid foods on which a child might choke (such as candy, hot dogs, peanuts, popcorn).

- Prevent Fires and Burns**
- Develop and practice a fire escape plan.
 - Check your smoke detector to make sure it is working.
 - Keep a fire extinguisher in or near the kitchen.
 - Check food temperatures carefully, especially if foods have been heated in a microwave oven.
 - Keep hot foods and liquids out of reach.
 - Put plastic covers in unused electrical outlets.
 - Throw away cracked or frayed old electrical cords.
 - Turn the water heater down to 120°F (50°C).

- Avoid Falls**
- Keep crib and playpen sides up.
 - Avoid using walkers.
 - Install safety gates to guard stairways.
 - Lock doors to dangerous areas like the basement or garage.
 - Check drawers, tall furniture, and lamps to make sure they can't fall over easily.

- Prevent Poisoning**
- Keep all medicines, vitamins, cleaning fluids, and gardening chemicals locked away or disposed of safely.
 - Install safety latches on cabinets.
 - Keep the poison center number on all phones. (1-800-222-1222)

Immunizations - At the 6-month visit, your baby will typically receive a:

- DTaP (diphtheria, tetanus, acellular pertussis) shot
- Hib (Haemophilus influenza type B) shot
- PCV-7 (Pneumococcal conjugate vaccine - 7) shot.

Your baby may run a fever and be irritable for about 1 day after the shots. Your baby may also have some soreness, redness, and swelling in the area where the shots were given. Acetaminophen drops (3/4 dropper, or 0.6 ml, every 4 to 6 hours) as needed for fever and irritability. For swelling or soreness, put a wet, warm washcloth on the area of the shots as often and as long as needed to provide comfort.

Call your child's physician if:

- Your child has a rash or any reaction to the shots other than fever and mild irritability.
- Your child has a fever that lasts more than 36 hours.

Next Visit - Your baby's next routine visit should be at the age of 9 months. Please bring the shot card to each visit.

